

EDITORIAL



Best practices in airway management in critically ill adults

Audrey De Jong^{1,2} , Samir Jaber^{1,2} and Sheila Nainan Myatra^{3*}

© 2025 Springer-Verlag GmbH Germany, part of Springer Nature

Tracheal intubation is one of the most common procedures performed in the critically ill. In a global study including 2964 critically ill adult patients, at least one major peri-intubation adverse event occurred in 45% of patients, with cardiovascular collapse (43%) and severe hypoxemia (9%) being the most common [1]. Of note, the occurrence of these events was associated with an increased risk of both intensive care unit (ICU) and 28-day mortality [1]. Factors contributing to this high complication rate include the complex environment, the varied experience of the operator, and most importantly, the critical illness itself [2, 3]. Critically ill patients have a *physiologically difficult airway*, defined in a recent global expert consensus Delphi study as one in which the patient's physiological and pathophysiological alterations increase the risk of complications during tracheal intubation and transition to positive pressure ventilation [4, 5]. Airway societies and experts have provided guidance to mitigate the complications that occur during this high-risk procedure [2–6]. This narrative review highlights various strategies to improve the safety and efficacy of tracheal intubation in critically ill adults. Though literature including critically ill patients was reviewed, some papers had mixed populations from ICU, operating room (OR), and emergency settings, which may make the extrapolations difficult between different settings. Future research priorities based on the gaps identified by the authors of this paper have been outlined in Table 1.

Preparation for tracheal intubation

The MACOCHA score [7], which considers physiological alterations and the experience of the operator, in addition to anatomical difficulty (Supplemental Table 1), is validated for airway assessment in ICU. Human factors influence safe performance of tracheal intubation, with team preparation being paramount to mitigate the risk of complications [5]. Though the benefit of using a pre-intubation checklist to reduce complications has not been established, this may be effective when interventions for physiological optimization are included [8] and when used by less experienced teams.

Respiratory optimization

The use of head-elevated laryngoscopic position (HELP), with the head of the bed elevated to 30 degrees, has been suggested by experts [5]. The posterior portions of the lung become more prone to atelectasis and collapse in the supine position. A head-up position during preoxygenation helps to increase the functional residual capacity (FRC) and thereby decrease the incidence of hypoxemia during tracheal intubation, especially in patients with obesity [5, 9].

Several preoxygenation strategies have been practiced to reduce the incidence of hypoxemia during tracheal intubation: bag-valve mask (self-inflating device), non-invasive ventilation (NIV) which provides positive pressure ventilation, non-rebreather mask (high-concentration oxygen delivery device), or high-flow nasal oxygen (HFNO). Preoxygenation using a facemask (standard oxygen or NIV) needs to be removed during laryngoscopy, while HFNO has the advantage that it can be continued. The PREOXI trial [10] confirms the superiority of NIV over facemask oxygen (either a nonrebreather mask or bag-valve mask device) for preoxygenation, without increased incidence of aspiration. No difference in severe hypoxemia was noted when preoxygenation with NIV or

*Correspondence: sheila150@hotmail.com

³ Department of Anaesthesiology, Critical Care and Pain, Tata Memorial Hospital, Homi Bhabha National Institute, Mumbai, India
Full author information is available at the end of the article

Table 1 Future research priorities for airway management in critically ill adults

1	Determine the optimal strategy to prevent cardiovascular collapse (using a fluid bolus, or vasopressors or a combination prior to tracheal intubation) during tracheal intubation
2	Determine which vasopressor (ephedrine, phenylephrine, norepinephrine) could be used and how (dose, timing, dilution) to prevent cardiovascular collapse during tracheal intubation
3	Compare the efficacy of NIV, HFNO, combinations of the two and other methods, in different patient populations, clinical settings, and levels of hypoxemia, including the evaluation of long-term effects
4	Evaluate the role of peri-procedural point-of-care ultrasound for the assessment and management of cardiorespiratory compromise during tracheal intubation in the critically ill
5	Evaluate the ideal combination of sedative-hypnotic drugs to enhance the efficacy and safety of tracheal intubation in the critically ill
6	Identify patients that could benefit from the Sellick maneuver performed during rapid sequence induction
7	Assess whether the use of videolaryngoscope compared to direct laryngoscope is associated with fewer episodes of hypoxemia and other complications, in addition to first pass intubation success
8	Assess the impact of individualized protocols for tracheal intubation based on the severity of illness of the patient
9	Assess the role of artificial intelligence to help to apply best practices for tracheal intubation in critically ill patients and assess its efficacy in randomized controlled trials
10	Establish how tracheal intubation in critically ill adults can be performed safely and effectively in resource limited setting

NIV non-invasive ventilation, HFNO high flow nasal oxygen

HFNO (continued during laryngoscopy) was compared in hypoxemic patients undergoing tracheal intubation. A trial in patients with acute hypoxemic respiratory failure comparing preoxygenation with NIV versus HFNO (continued during laryngoscopy) did not show any difference in the incidence of severe hypoxemia. However, the subgroup analysis suggested that NIV may be superior to HFNO among patients with moderate to severe hypoxemia [11]. A proof of concept study showed that NIV combined with HFNO was superior to NIV alone in preventing desaturation following intubation of hypoxemic patients in ICU. This needs further evaluation [12]. A systematic review comparing preoxygenation strategies in critically ill adults concluded that NIV or HFNO are superior to facemask oxygen during tracheal intubation and that compared with HFNO, NIV decreases the incidence of hypoxemia during tracheal intubation [13]. These findings suggest that it may be time to abandon the use of conventional facemask oxygen in this high-risk population [14]. In agitated patients, the use of a sub-anesthetic dose of ketamine to facilitate preoxygenation (i.e., delayed sequence intubation) significantly decreases peri-intubation hypoxia [15]. Gentle mask ventilation performed during the apneic phase of a modified rapid sequence intubation (RSI) has been shown to reduce peri-procedural hypoxemia without increasing the incidence of aspiration [16]. Research gaps in the field of respiratory optimization are detailed in Table 1.

Hemodynamic optimization

Considering the high incidence of hemodynamic collapse during the procedure, hemodynamic optimization and monitoring are paramount. However, the precise method of performing hemodynamic optimization is not clear in

the literature, and further research is needed (Table 1). The Montpellier intubation protocol, the use of which demonstrated a reduction in intubation-related complications, includes a fluid bolus prior to tracheal intubation, unless contraindicated [8]. This has been validated by another team in an observational study [17]. However, in two randomized trials, administration of a fluid bolus prior to intubation showed no reduction in cardiovascular collapse [18, 19]. Two ongoing international trials (the FLUVA Trial [NCT05318066] and the PREVENTION trial [NCT05014581]) are investigating the effectiveness of pre-emptively administering vasopressors in preventing cardiovascular collapse in critically ill adults undergoing tracheal intubation.

Performing tracheal intubation

A modified RSI technique (administration of rapid-onset sedative hypnotic and a rapid-acting neuromuscular blocking agent, and/or gentle mask ventilation) should be considered to minimize the risk of aspiration [5]. Cricoid pressure (if applied) should be pursued until the tracheal tube is inserted appropriately and the tracheal tube cuff is inflated, as regurgitation may occur with the pressure transmitted to the esophagus. However, it should be released in case of difficulty in glottic visualization [5]. Evidence is lacking to determine which patients could benefit from the use of cricoid pressure (Table 1).

Drugs used for tracheal intubation

The post hoc analysis of the INTUBE study showed that the use of propofol for induction was the only modifiable independent predictor of cardiovascular collapse in these patients [20]. Therefore, drugs with a more stable hemodynamic profile, such as ketamine and etomidate, have

been recommended [5]. The decision to use either of these drugs should be based on individual patient characteristics, contra-indications, side effects of each drug, and local practices [5]. Future studies are required to individualize induction agents used for tracheal intubation in the critically ill (Table 1).

Use of neuromuscular blockade during RSI is associated with greater first-pass intubation success and lesser complications. A study comparing succinylcholine and rocuronium for tracheal intubation in critically ill patients showed no clinically meaningful difference between the two drugs [21]. Succinylcholine should be avoided in patients with skeletal muscle myopathies, known allergy, history of malignant hyperthermia, hyperkalemia, and significant burn injury [5]. There is a dose-response effect of rocuronium; higher doses of at least 1.2 mg/kg are required to produce faster time to optimal intubating conditions [2]. Sugammadex may be used for the rapid reversal of rocuronium in a “cannot intubate, cannot ventilate” situation, though the safety data in critically ill patients is limited [5]. However, its role in this situation is extremely limited, as sugammadex may reverse the paralytic, but not the need for intubation, required in most of these patients [22].

Devices used for tracheal intubation

Various tools and strategies can help improve first-pass intubation success and thereby reduce complications. It is now clearly demonstrated that videolaryngoscopy (VL) provides superior intubation conditions over direct laryngoscopy (DL) [23, 24]. The STYLETO (Stylet for orotracheal intubation) trial demonstrated that using a stylet with a tracheal tube during intubation with a DL resulted in a significantly higher first-pass intubation success rate compared to a tracheal tube alone [25]. A study in critically ill adults comparing the use of a tracheal tube with a bougie or a stylet during the first attempt at intubation reported no difference in first-pass intubation success [26]. Taken together, these studies highlight the importance of the routine use of a stylet or a bougie during the first attempt at tracheal intubation. Nevertheless, future research is needed to assess which of these devices may be superior in specific settings (Table 1).

Post procedure care

Tracheal placement of the tube should be confirmed using consistent waveform capnography over at least seven breaths [24]. Chest auscultation, chest X-ray, and/or bronchoscopy are acceptable methods to confirm the optimal depth of the tube in the trachea [5]. A lung protective ventilation strategy, along with the use of sedative infusions, should be employed post intubation [5]. Complications that occur should be managed appropriately.

Complications associated with tracheal intubation in critically ill adults continue to remain high [1]. While evidence supporting the management of patients with a *physiologically difficult airway* is evolving, the implementation of the existing strategies discussed above may help improve patient outcomes.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1007/s00134-025-08181-9>.

Author details

¹ Department of Anesthesiology and Critical Care Medicine B (DAR B), Saint-Eloi Hospital, University Teaching Hospital of Montpellier, Montpellier, France. ² INSERM U1046 PhyMedExp, Montpellier, France. ³ Department of Anaesthesiology, Critical Care and Pain, Tata Memorial Hospital, Homi Bhabha National Institute, Mumbai, India.

Funding

Nil.

Data availability

Not applicable.

Declarations

Conflict of interest

ADJ has received payment/honoraria from Medtronic, Sanofi, Drager, Fisher-Paykel, and Viatrix for lectures, presentations, speakers' bureaus, manuscript writing or educational events. SJ has received consulting fees from Drager, Medtronic, Mindray, Fresenius, Baxter, and Fisher & Paykel. SNM declares no conflict of interest.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 28 June 2025 Accepted: 21 October 2025

Published: 3 November 2025

References

- Russotto V, Myatra SN, Laffey JG, Tassistro E, Antolini L, Bauer P, Lascarrou JB, Szuldrzynski K, Camporota L, Pelosi P, Sorbello M, Higgs A, Greif R, Putensen C, Agvald-Öhman C, Chalkias A, Bokums K, Brewster D, Rossi E, Fumagalli R, Pesenti A, Foti G, Bellani G (2021) Intubation practices and adverse peri-intubation events in critically ill patients from 29 countries. *JAMA* 325:1164–1172
- Myatra SN, Ahmed SM, Kundra P, Garg R, Ramkumar V, Patwa A, Shah A, Raveendra US, Shetty SR, Doctor JR, Pawar DK, Ramesh S, Das S, Divatia JV (2017) Republication: All India Difficult Airway Association 2016 guidelines for tracheal intubation in the intensive care unit. *Indian J Crit Care Med* 21:146–153
- Higgs A, McGrath BA, Goddard C, Rangasami J, Suntharalingam G, Gale R, Cook TM (2018) Guidelines for the management of tracheal intubation in critically ill adults. *Br J Anaesth* 120:323–352
- Myatra SN, Divatia JV, Brewster DJ (2022) The physiologically difficult airway: an emerging concept. *Curr Opin Anaesthesiol* 35:115–121
- Karamchandani K, Nasa P, Jarzebowski M, Brewster DJ, De Jong A, Bauer PR, Berkow L, Brown CA 3rd, Cabrini L, Casey J, Cook T, Divatia JV, Duggan LV, Ellard L, Ergon B, JonssonFagerlund M, Gatward J, Greif R, Higgs A, Jaber S, Janz D, Joffe AM, Jung B, Kovacs G, Kwizera A, Laffey JG, Lascarrou JB, Law JA, Marshall S, McGrath BA, Mosier JM, Perin D, Roca O, Rollé A, Russotto V, Sakles JC, Shrestha GS, Smischney NJ, Sorbello M, Tung A,

- Jabaley CS, Myatra SN (2024) Tracheal intubation in critically ill adults with a physiologically difficult airway. An international Delphi study. *Intensive Care Med* 50:1563–1579
6. De Jong A, Myatra SN, Roca O, Jaber S (2022) How to improve intubation in the intensive care unit. Update on knowledge and devices. *Intensive Care Med* 48:1287–1298
 7. De Jong A, Molinari N, Terzi N, Mongardon N, Arnal J-M, Guitton C, Allaouchiche B, Paugam-Burtz C, Constantin J-M, Lefrant J-Y (2013) Early identification of patients at risk for difficult intubation in the intensive care unit: development and validation of the MACOCHA score in a multicenter cohort study. *Am J Respir Crit Care Med* 187:832–839
 8. Jaber S, Jung B, Corne P, Sebbane M, Muller L, Chanques G, Verzilli D, Jonquet O, Eledjam J-J, Lefrant J-Y (2010) An intervention to decrease complications related to endotracheal intubation in the intensive care unit: a prospective, multiple-center study. *Intensive Care Med* 36:248–255
 9. Dixon BJ, Dixon JB, Carden JR, Burn AJ, Schachter LM, Playfair JM, Laurie CP, O'Brien PE (2005) Preoxygenation is more effective in the 25 degrees head-up position than in the supine position in severely obese patients: a randomized controlled study. *Anesthesiology* 102:1110–1115 (**discussion 1115A**)
 10. Gibbs KW, Semler MW, Driver BE, Seitz KP, Stemppek SB, Taylor C, Resnick-Ault D, White HD, Gandotra S, Doerschug KC, Mohamed A, Prekker ME, Khan A, Gaillard JP, Andrea L, Aggarwal NR, Brainard JC, Barnett LH, Halliday SJ, Blinder V, Dagan A, Whitson MR, Schauer SG, Walker JE Jr, Barker AB, Palakshappa JA, Muhs A, Wozniak JM, Kramer PJ, Withers C, Ghamande SA, Russell DW, Schwartz A, Moskowitz A, Hansen SJ, Allada G, Goranson JK, Fein DG, Sottile PD, Kelly N, Alwood SM, Long MT, Malhotra R, Shapiro NI, Page DB, Long BJ, Thomas CB, Trent SA, Janz DR, Rice TW, Self WH, Bebartta VS, Lloyd BD, Rhoads J, Womack K, Imhoff B, Ginde AA, Casey JD (2024) Noninvasive ventilation for preoxygenation during emergency intubation. *N Engl J Med* 390:2165–2177
 11. Frat JP, Ricard JD, Quenot JP, Pichon N, Demoule A, Forel JM, Mira JP, Coudroy R, Berquier G, Voisin B, Colin G, Pons B, Danin PE, Devaquet J, Prat G, Clere-Jehl R, Petitpas F, Vivier E, Razazi K, Nay MA, Souday V, Delamonica J, Argaud L, Ehrmann S, Gibelin A, Girault C, Andreu P, Vignon P, Dangers L, Ragot S, Thille AW (2019) Non-invasive ventilation versus high-flow nasal cannula oxygen therapy with apnoeic oxygenation for preoxygenation before intubation of patients with acute hypoxaemic respiratory failure: a randomised, multicentre, open-label trial. *Lancet Respir Med* 7:303–312
 12. Jaber S, Monnin M, Girard M, Conseil M, Cisse M, Carr J, Mahul M, Delay JM, Belafia F, Chanques G (2016) Apnoeic oxygenation via high-flow nasal cannula oxygen combined with non-invasive ventilation preoxygenation for intubation in hypoxaemic patients in the intensive care unit: the single-centre, blinded, randomised controlled OPTINIV trial. *Intensive Care Med* 42:1877–1887
 13. Pitre T, Liu W, Zeraatkar D, Casey JD, Dionne JC, Gibbs KW, Ginde AA, Needham-Nethercott N, Rice TW, Semler MW, Rochweg B (2025) Preoxygenation strategies for intubation of patients who are critically ill: a systematic review and network meta-analysis of randomised trials. *Lancet Respir Med* 13:585–596
 14. Myatra SN, Jaber S (2025) Preoxygenation for tracheal intubation in adults who are critically ill: time to abandon the facemask? *Lancet Respir Med* 13:570–571
 15. Bandyopadhyay A, Kumar P, Jafra A, Thakur H, Yaddanapudi LN, Jain K (2023) Peri-intubation hypoxia after delayed versus rapid sequence intubation in critically injured patients on arrival to trauma triage: a randomized controlled trial. *Anesth Analg* 136:913–919
 16. Casey JD, Janz DR, Russell DW, Vonderhaar DJ, Joffe AM, Dischert KM, Brown RM, Zouk AN, Gulati S, Heideman BE, Lester MG, Toporek AH, Bentov I, Self WH, Rice TW, Semler MW (2019) Bag-mask ventilation during tracheal intubation of critically ill adults. *N Engl J Med* 380:811–821
 17. Corl KA, Dado C, Agarwal A, Azab N, Amass T, Marks SJ, Levy MM, Merchant RC, Aliotta J (2018) A modified Montpellier protocol for intubating intensive care unit patients is associated with an increase in first-pass intubation success and fewer complications. *J Crit Care* 44:191–195
 18. Janz DR, Casey JD, Semler MW, Russell DW, Dargin J, Vonderhaar DJ, Dischert KM, West JR, Stemppek S, Wozniak J, Caputo N, Heideman BE, Zouk AN, Gulati S, Stigler WS, Bentov I, Joffe AM, Rice TW (2019) Effect of a fluid bolus on cardiovascular collapse among critically ill adults undergoing tracheal intubation (PrePARE): a randomised controlled trial. *Lancet Respir Med* 7:1039–1047
 19. Russell DW, Casey JD, Gibbs KW, Ghamande S, Dargin JM, Vonderhaar DJ, Joffe AM, Khan A, Prekker ME, Brewer JM, Dutta S, Landsperger JS, White HD, Robison SW, Wozniak JM, Stemppek S, Barnes CR, Krol OF, Arroliga AC, Lat T, Gandotra S, Gulati S, Bentov I, Walters AM, Dischert KM, Nonas S, Driver BE, Wang L, Lindsell CJ, Self WH, Rice TW, Janz DR, Semler MW (2022) Effect of fluid bolus administration on cardiovascular collapse among critically ill patients undergoing tracheal intubation: a randomized clinical trial. *JAMA* 328:270–279
 20. Russotto V, Tassistro E, Myatra SN, Parotto M, Antolini L, Bauer P, Lascarrou JB, Szuldrzyński K, Camporota L, Putensen C, Pelosi P, Sorbello M, Higgs A, Greif R, Pesenti A, Valsecchi MG, Fumagalli R, Foti G, Bellani G, Laffey JG (2022) Peri-intubation cardiovascular collapse in patients who are critically ill: insights from the INTUBE study. *Am J Respir Crit Care Med* 206:449–458
 21. Guihard B, Chollet-Xémar C, Lakhnati P, Vivien B, Broche C, Savary D, Ricard-Hibon A, Marianne DiCassou PJ, Adnet F, Wiel E, Deutsch J, Tissier C, Loeb T, Bounes V, Rousseau E, Jabre P, Huiart L, Ferdynus C, Combes X (2019) Effect of rocuronium vs succinylcholine on endotracheal intubation success rate among patients undergoing out-of-hospital rapid sequence intubation: a randomized clinical trial. *JAMA* 322:2303–2312
 22. Jansson PS, Pimentel MPT, Seethala RR (2024) Emergency department usage of sugammadex in a large regional health system. *Acad Emerg Med* 31:713–715
 23. Prekker ME, Driver BE, Trent SA, Resnick-Ault D, Seitz KP, Russell DW, Gaillard JP, Latimer AJ, Ghamande SA, Gibbs KW, Vonderhaar DJ, Whitson MR, Barnes CR, Walco JP, Douglas IS, Krishnamoorthy V, Dagan A, Bastman JJ, Lloyd BD, Gandotra S, Goranson JK, Mitchell SH, White HD, Palakshappa JA, Espinera A, Page DB, Joffe A, Hansen SJ, Hughes CG, George T, Herbert JT, Shapiro NI, Schauer SG, Long BJ, Imhoff B, Wang L, Rhoads JP, Womack KN, Janz DR, Self WH, Rice TW, Ginde AA, Casey JD, Semler MW (2023) Video versus direct laryngoscopy for tracheal intubation of critically ill adults. *N Engl J Med* 389:418–429
 24. Russotto V, Lascarrou JB, Tassistro E, Parotto M, Antolini L, Bauer P, Szuldrzyński K, Camporota L, Putensen C, Pelosi P, Sorbello M, Higgs A, Greif R, Grasselli G, Valsecchi MG, Fumagalli R, Foti G, Caironi P, Bellani G, Laffey JG, Myatra SN (2023) Efficacy and adverse events profile of videolaryngoscopy in critically ill patients: subanalysis of the INTUBE study. *Br J Anaesth* 131:607–616
 25. Jaber S, Rollé A, Godet T, Terzi N, Riu B, Asfar P, Bourenne J, Ramin S, Lemiale V, Quenot J-P (2021) Effect of the use of an endotracheal tube and stylet versus an endotracheal tube alone on first-attempt intubation success: a multicentre, randomised clinical trial in 999 patients. *Intensive Care Med* 47:653–664
 26. Driver BE, Semler MW, Self WH, Ginde AA, Trent SA, Gandotra S, Smith LM, Page DB, Vonderhaar DJ, West JR, Joffe AM, Mitchell SH, Doerschug KC, Hughes CG, High K, Landsperger JS, Jackson KE, Howell MP, Robison SW, Gaillard JP, Whitson MR, Barnes CM, Latimer AJ, Koppurapu VS, Alvis BD, Russell DW, Gibbs KW, Wang L, Lindsell CJ, Janz DR, Rice TW, Prekker ME, Casey JD (2021) Effect of use of a bougie vs endotracheal tube with stylet on successful intubation on the first attempt among critically ill patients undergoing tracheal intubation: a randomized clinical trial. *JAMA* 326:2488–2497